



Serving Antrim, Charlevoix, Emmet & Otsego Counties

MEDIA CONTACT:
Erin Luckhardt, PIO
Office: 231-347-5628
Information Line: 1-800-386-5959
eluckhardt@nwhealth.org
nwhealth.org

For Immediate Release
March 4, 2020

Health Department of Northwest Michigan Advises Residents to Prepare not Panic for Coronavirus Disease 2019 (COVID-19)

The Health Department of Northwest Michigan is continuously working on preparedness and response planning for Coronavirus Disease 2019 (COVID-19). Communication occurs daily with federal, state and local officials/partners. The Health Department is monitoring the situation at a local level and coordinating efforts with other local health departments -- this includes Benzie-Leelanau District Health Department, Central Michigan District Health Department, District Health Department #2, District Health Department #4, District Health Department #10 and Grand Traverse County Health Department. Michigan, including Antrim, Charlevoix, Emmet and Otsego Counties have no confirmed cases at this time.

As COVID-19 cases increase worldwide, including in the United States, it is important to be mindful of preparing for a shift from an epidemic to a pandemic. At this time, we are encouraging our communities to *prepare, not panic*. It is still uncertain when COVID-19 will impact the greater United States, for how long, or the severity, and information is changing on a daily basis.

Health Officer, Lisa Peacock, reminds residents that "It is important to consider actions to reduce the spread of illness in general – things like staying home if you are ill, coughing into an elbow or tissue, and frequent handwashing are important. While the situation is regularly changing, it is also important to stay up to date with reliable sources and to begin identifying some actions that can help protect your health and those you care about."

These measures will help decrease and slow the spread of infection, and reduce the impact on our community:

- Pay attention, and comply, with public health requests and recommendations.
- Get your information from credible sources (Center for Disease Control and Prevention, Michigan Department of Health and Human Services, Health Department, etc.).
- Support your family, friends and neighbors when they need your help.
- Practice good hand washing hygiene – wash hands regularly or use an alcohol-based hand sanitizer (60% or more) if soap and water are not available.
- Stay home when you are sick (work, school, social gatherings, events, etc.).
- Clean all "high touch" surfaces every day with household cleaning spray or wipes.
- Cover coughs, sneezes and avoid touching your face.
- Avoid sharing personal household items.
- Be current on your routine vaccinations.
- Call ahead before going to the doctor or hospital if you are sick (most Coronavirus infections are not serious, but as always, be mindful of concerning symptoms like shortness of breath and prolonged fever).

- For additional information about Coronavirus Disease 2019, please visit

For information about Coronavirus Disease 2019 precautions for businesses, please visit <https://www.cdc.gov/coronavirus/2019-ncov/specific-groups/guidance-business-response.html>.

###



GRETCHEN WHITMER
GOVERNOR

STATE OF MICHIGAN
OFFICE OF THE GOVERNOR
LANSING

GARLIN GILCHRIST II
LT. GOVERNOR

Michigan Department of Health and Human Services (MDHHS)
Interim Recommendations for COVID-19 Community Mitigation Strategies
March 11, 2020

[The most up-to-date guidance on these and other mitigation strategies is available at www.Michigan.gov/coronavirus. This matter is rapidly evolving and MDHHS may provide updated guidance.]

Community mitigation strategies are crucial to slowing the transmission of coronavirus disease 2019 (COVID-19) in Michigan, particularly before a vaccine or treatment becomes available. These strategies provide essential protections to individuals at risk of severe illness and to health care and other critical infrastructure workforces. Preventing a sudden, sharp increase in the number of people infected with COVID-19 will help minimize disruptions to daily life and limit the demand on health care providers and facilities.

These recommended strategies apply at the individual, organizational, and community levels. They apply to businesses, workplaces, schools, community organizations, health care institutions, and individuals of all ages, backgrounds, and health profiles. Everyone has some measure of responsibility to help limit the spread of this disease. Even individuals who are healthy can help prevent the spread of COVID-19 to others.

Michiganders have been preparing for COVID-19 for weeks, and all individuals should continue to take the following basic personal-hygiene measures to prevent the spread of the virus:

- wash your hands often with soap and water or use hand sanitizer;
- avoid touching your eyes, nose, or mouth with unwashed hands;
- cover your mouth and nose with a tissue when coughing or sneezing;
- avoid handshakes;
- avoid contact with sick people who are sick; and
- stay home when you are sick.

Michigan must take further action, however. To help avoid a rapid increase of cases in the state, MDHHS recommends implementing now the following community mitigation strategies. Please note, guidance for health care providers is not addressed in this document, but is available at www.Michigan.gov/coronavirus.

Individuals and families at home:

1. Learn about the signs and symptoms of COVID-19. Symptoms include fever, cough, and difficulty breathing.
2. If you have respiratory symptoms, **STAY HOME WHEN YOU ARE SICK**. Call your health care provider's office in advance of your visit.
3. Regularly clean and disinfect frequently touched surfaces, like doorknobs, keyboards, cell phones, and light switches.
4. Communicate and reinforce best practices for washing hands and covering coughs and sneezes.
5. Be prepared if there is COVID-19 in your household or a disruption of daily activities in your community. For example, maintain a supply of medications, food, and other essentials in your house. Consider alternative shopping options such as curbside pickups or online deliveries.
6. Access services as much as possible online or by phone.

Individuals at risk of severe illness:

These individuals include, but are not limited to, older adults and persons of any age with underlying medical conditions, such as persons with a blood disorder (e.g., sickle cell disease or a disorder being treated with blood thinners), an endocrine disorder (e.g., diabetes mellitus), or a metabolic disorder (such as inborn error of metabolism); those with heart disease, lung disease (including asthma or chronic obstructive pulmonary disease), chronic kidney disease, or chronic liver disease; those with a compromised immune system (e.g., those who are receiving treatments such as radiation or chemotherapy, who have received an organ or bone marrow transplant, who are taking high doses of immunosuppressant, or who have HIV or AIDS); those who are currently pregnant or were pregnant in the last two weeks; and those with neurological or neurologic and neurodevelopment conditions.

1. Individuals at risk of severe illness should stay at home and keep away from others who are sick, except in exceptional circumstances. Wash your hands often, particularly after contact with high-touch surfaces. Avoid crowds and closed-in settings with little air ventilation as much as possible. Avoid cruise travel and non-essential air travel.
2. Regularly clean and disinfect frequently touched surfaces, like doorknobs, keyboards, cell phones, and light switches.
3. In households with individuals at risk of severe illness, provide a protected space for those individuals and have healthy people conduct themselves as if they were a significant risk to those individuals. For example, healthy people should wash their hands before feeding or caring for an at-risk individual.
4. Have a plan for if you get sick, and stay in touch with others by phone or email.
5. Watch for symptoms and emergency warning signs that require immediate medical attention.
6. Family members and caregivers can support older adults by knowing what medications they are taking and ensuring there is an extra supply on hand.
7. Family members and caregivers can support older adults by monitoring food and other necessary medical supplies (e.g., oxygen, incontinence, dialysis, and wound care supplies) and by creating a back-up plan for securing those essentials if they run out.

Assisted living facilities, adult day care programs, and senior living facilities:

1. Encourage personal protective measures among residents, clients, and staff, including handwashing and covering coughs and sneezes.
2. Encourage residents, clients, and staff to stay home (or in their private quarters) when sick and to notify program administrators of illness.
3. Regularly clean and disinfect frequently touched surfaces, like doorknobs, keyboards, cell phones, and light switches.
4. Ensure hand hygiene supplies are readily accessible throughout the facility.
5. If caring for an individual living in a care facility, monitor the situation, ask about the health of the other residents frequently, and know the protocol if there is an outbreak.
6. Implement social distancing measures as feasible, such as reducing large gatherings, altering meal schedules to reduce mixing, and limiting programs with external staff.

7. Limit visitors to only those who are absolutely necessary and implement screening of visitors for temperature and respiratory symptoms.
8. Maintain contact with individuals at risk of severe illness who are no longer able to attend day care programs.

Universities and colleges:

1. Educate students and the community about COVID-19 and preventative hygiene practices.
2. Regularly clean and disinfect frequently touched surfaces, like doorknobs, keyboards, cell phones, and light switches.
3. Encourage staff and students to stay home when sick.
4. Cancel or postpone large gatherings, conferences, and sporting events (e.g., greater than 100 people in a shared space).
5. Consider tele-learning opportunities, as feasible.
6. Communicate and reinforce best practices for washing hands and covering coughs and sneezes.
7. Students and faculty at risk of severe illness should implement plans for distance learning.

Schools and childcare facilities:

1. Educate students and the community about COVID-19 and preventative hygiene practices.
2. Encourage staff and students to stay home when sick.
3. Report influenza-like activity, absenteeism, and potential school dismissals to public health officials.
4. Separate sick students and staff from others until they can go home. When feasible, identify a “sick room” through which others do not regularly pass.
5. Communicate and reinforce best practices for washing hands and covering coughs and sneezes.

6. Regularly clean and disinfect frequently touched surfaces, like doorknobs, keyboards, cell phones, and light switches.
7. Consider having students eat meals in their classrooms as opposed to a cafeteria-like setting, or alter meal schedules for smaller-group gatherings.
8. Limit large gatherings such as assemblies and limit inter-school interactions.
9. Alter schedules for recess and entry/dismissal to reduce the mixing of large numbers of students and staff.
10. Schools with students at risk of severe illness should consider implementing plans for distance learning for those students, to the extent feasible.

Workplaces:

1. Encourage employees to stay home when sick and to notify supervisors of illness.
2. Communicate and reinforce best practices for washing hands covering coughs and sneezes.
3. Regularly clean and disinfect frequently touched surfaces, like doorknobs, keyboards, cell phones, and light switches.
4. Ensure hand hygiene supplies are readily accessible throughout the workplace.
5. Encourage staff to tele-work when feasible, particularly individuals at risk of severe illness.
6. Implement social distancing measures as feasible, including limiting in-person meetings.
7. Limit large work-related gatherings (e.g., staff meetings and after-work functions).
8. Limit non-essential work travel.
9. Cancel or postpone large gatherings, conferences, and sporting events (e.g., greater than 100 people in a shared space).
10. Discourage employees from eating meals in a large group setting, such as a cafeteria.
11. Tailor continuity of operation plans to the COVID-19 threat.

Community and faith-based organizations:

1. Identify safe ways to serve those who are at high risk or vulnerable through outreach and assistance.
2. Encourage staff and members to stay home when sick and to notify the organization of illness.
3. Communicate and reinforce best practices for washing hands and covering coughs and sneezes.
4. Regularly clean and disinfect frequently touched surfaces, like doorknobs, keyboards, cell phones, and light switches.
5. Ensure hand hygiene supplies are readily accessible.
6. Implement social distancing measures as feasible.
7. Reduce in-person gatherings and activities, especially for organizations with individuals at risk of severe illness. Consider offering video or audio of events.
8. Determine ways to continue providing support services to individuals at risk of severe illness while limiting group settings and exposures.
9. Avoid large gatherings (e.g., greater than 100 people in a shared space) or move to smaller and staggered gatherings.
10. For organizations that serve high-risk communities, cancel gatherings of more than 10 people.

Other mass events:

1. Cancel or postpone large gatherings, conferences, and sporting events (e.g., greater than 100 people in a shared space).
2. Regularly clean and disinfect frequently touched surfaces, like doorknobs, keyboards, and light switches.
3. Communicate and reinforce best practices for washing hands and covering coughs and sneezes.



What to do if you are sick with coronavirus disease 2019 (COVID-19)

If you are sick with COVID-19 or suspect you are infected with the virus that causes COVID-19, follow the steps below to help prevent the disease from spreading to people in your home and community.

Stay home except to get medical care

You should restrict activities outside your home, except for getting medical care. Do not go to work, school, or public areas. Avoid using public transportation, ride-sharing, or taxis.

Separate yourself from other people and animals in your home

People: As much as possible, you should stay in a specific room and away from other people in your home. Also, you should use a separate bathroom, if available.

Animals: Do not handle pets or other animals while sick. See [COVID-19 and Animals](#) for more information.

Call ahead before visiting your doctor

If you have a medical appointment, call the healthcare provider and tell them that you have or may have COVID-19. This will help the healthcare provider's office take steps to keep other people from getting infected or exposed.

Wear a facemask

You should wear a facemask when you are around other people (e.g., sharing a room or vehicle) or pets and before you enter a healthcare provider's office. If you are not able to wear a facemask (for example, because it causes trouble breathing), then people who live with you should not stay in the same room with you, or they should wear a facemask if they enter your room.

Cover your coughs and sneezes

Cover your mouth and nose with a tissue when you cough or sneeze. Throw used tissues in a lined trash can; immediately wash your hands with soap and water for at least 20 seconds or clean your hands with an alcohol-based hand sanitizer that contains at least 60% alcohol covering all surfaces of your hands and rubbing them together until they feel dry. Soap and water should be used preferentially if hands are visibly dirty.

Avoid sharing personal household items

You should not share dishes, drinking glasses, cups, eating utensils, towels, or bedding with other people or pets in your home. After using these items, they should be washed thoroughly with soap and water.

Clean your hands often

Wash your hands often with soap and water for at least 20 seconds. If soap and water are not available, clean your hands with an alcohol-based hand sanitizer that contains at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry. Soap and water should be used preferentially if hands are visibly dirty. Avoid touching your eyes, nose, and mouth with unwashed hands.

Clean all "high-touch" surfaces every day

High touch surfaces include counters, tabletops, doorknobs, bathroom fixtures, toilets, phones, keyboards, tablets, and bedside tables. Also, clean any surfaces that may have blood, stool, or body fluids on them. Use a household cleaning spray or wipe, according to the label instructions. Labels contain instructions for safe and effective use of the cleaning product including precautions you should take when applying the product, such as wearing gloves and making sure you have good ventilation during use of the product.

Monitor your symptoms

Seek prompt medical attention if your illness is worsening (e.g., difficulty breathing). **Before** seeking care, call your healthcare provider and tell them that you have, or are being evaluated for, COVID-19. Put on a facemask before you enter the facility. These steps will help the healthcare provider's office to keep other people in the office or waiting room from getting infected or exposed.

Ask your healthcare provider to call the local or state health department. Persons who are placed under active monitoring or facilitated self-monitoring should follow instructions provided by their local health department or occupational health professionals, as appropriate. When working with your local health department check their available hours.

If you have a medical emergency and need to call 911, notify the dispatch personnel that you have, or are being evaluated for COVID-19. If possible, put on a facemask before emergency medical services arrive.

Discontinuing home isolation

Patients with confirmed COVID-19 should remain under home isolation precautions until the risk of secondary transmission to others is thought to be low. The decision to discontinue home isolation precautions should be made on a case-by-case basis, in consultation with healthcare providers and state and local health departments.



For more information: www.cdc.gov/COVID19